

Scotland's Rural College

Co-infection with *Fasciola hepatica* may increase the risk of *Escherichia coli* O157 shedding in British cattle destined for the food chain

Howell, AK; Tongue, SC; Currie, C; Evans, J; Williams, DJL; McNeilly, TN

Published in:
Preventive Veterinary Medicine

DOI:
[10.1016/j.prevetmed.2017.12.007](https://doi.org/10.1016/j.prevetmed.2017.12.007)

First published: 07/12/2017

Document Version
Peer reviewed version

[Link to publication](#)

Citation for pulished version (APA):

Howell, AK., Tongue, SC., Currie, C., Evans, J., Williams, DJL., & McNeilly, TN. (2017). Co-infection with *Fasciola hepatica* may increase the risk of *Escherichia coli* O157 shedding in British cattle destined for the food chain. *Preventive Veterinary Medicine*, 150, 70 - 76. <https://doi.org/10.1016/j.prevetmed.2017.12.007>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Co-infection with *Fasciola hepatica* may increase the**
2 **risk of *Escherichia coli* O157 shedding in British cattle**
3 **destined for the food chain**

4 Alison K. Howell^{a*}, Sue C. Tongue^b, Carol Currie^c, Judith Evans^b, Diana J. L. Williams^{a,d**} and
5 Tom N. McNeilly^{c**}

6 ^aDepartment of Infection Biology, Institute of Infection and Global Health, University of
7 Liverpool, Liverpool, L69 7ZJ, UK

8 ^bSRUC Research, Epidemiology Research Unit, (Inverness Campus), Scotland's Rural College
9 (SRUC), Kings Buildings, West Mains Road, Edinburgh, EH9 3JG, U.K.

10 ^cMoredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik, Midlothian,
11 EH26 0PZ, UK

12 ^dSchool of Veterinary Science, University of Liverpool, Leahurst, Chester High Road, Neston,
13 CH64 7TE, UK.

14 *Corresponding author. Tel: +44 151 795 0235. E-mail address:

15 alison.howell@liverpool.ac.uk

16 **Joint senior authors

17 **Abstract**

18 *Escherichia coli* O157 is a zoonotic bacterium that can cause haemorrhagic diarrhoea in
19 humans and is of worldwide public health concern. Cattle are considered to be the main
20 reservoir for human infection. *Fasciola hepatica* is a globally important parasite of ruminant

21 livestock that is known to modulate its host's immune response and affect susceptibility to
22 bacterial pathogens such as *Salmonella* Dublin. Shedding of *E. coli* O157 is triggered by
23 unknown events, but the immune system is thought to play a part. We investigated the
24 hypothesis that shedding of *E. coli* O157 is associated with *F. hepatica* infection in cattle.
25 Three hundred and thirty four cattle destined for the food chain, from 14 British farms,
26 were tested between January and October 2015. *E. coli* O157 was detected by
27 immunomagnetic separation and bacterial load enumerated. *F. hepatica* infection status
28 was assessed by copro-antigen ELISA. A significant association ($p = 0.01$) was found
29 between the log percent positivity (PP) of the *F. hepatica* copro-antigen ELISA and *E. coli*
30 O157 shedding when the fixed effects of day of sampling and the age of the youngest
31 animal in the group, plus the random effect of farm were adjusted for. The results should
32 be interpreted cautiously due to the lower than predicted level of fluke infection in the
33 animals sampled. Nevertheless these results indicate that control of *F. hepatica* infection
34 may have an impact on the shedding of *E. coli* O157 in cattle destined for the human food
35 chain.

36 **Keywords: Escherichia coli O157, Fasciola hepatica, cattle, co-infection**

37 **1 Introduction**

38 *Fasciola hepatica*, or the common liver fluke, is a parasite of ruminant livestock, occurring
39 worldwide. Various studies have shown that *F. hepatica* can affect host immunity to other
40 pathogens (Moreau and Chauvin, 2010), by making the host more susceptible to infection
41 (Aitken et al., 1979, 1981; Brady et al., 1999); changing the pathogenesis of disease (Garza-
42 Cuartero et al., 2016); and interfering with diagnostic tests (DEFRA, 2005; Flynn et al.,
43 2007). This happens because infection with *F. hepatica* induces a mixed T helper type-2
44 (Th2) and T-regulatory response, with increased production of IL4, IL5, IL10, IL13 and TGF β ,

45 whilst T helper type-1 (Th1) responses are down regulated (Flynn et al., 2010; Graham-
46 Brown, 2016).

47 *Escherichia coli* O157 is a zoonotic bacterium that occurs worldwide and can cause
48 haemorrhagic diarrhoea in humans as a result of systemic Shiga toxin (Stx) activity. Cattle
49 are considered the main source of human infection, either through direct contact or
50 through contaminated food (Locking et al., 2001; Strachan et al., 2006; Gyles, 2007). An
51 estimated 20-40% of British cattle herds are reported to shed *E. coli* O157 (Paiba et al.,
52 2003; Ellis-Iversen et al., 2007; Gunn et al., 2007; Pearce et al., 2009). The annual reported
53 incidence of human *E. coli* O157 is 1.8 culture positive cases per 100,000 population in
54 England and Wales, and 4.5 cases per 100,000 in Scotland (Health Protection Network,
55 2013; Public Health England, 2013). In a proportion of cases, mainly in young children,
56 haemolytic uraemic syndrome may occur, and is potentially fatal (Chase-Topping et al.,
57 2008).

58 *Escherichia coli* O157 infections in cattle are usually asymptomatic as cattle lack vascular
59 receptors for Stx (Pruimboom-Brees et al., 2000), but both cellular and humoral immune
60 responses are induced and are required for immunity to *E. coli* O157 (Corbishley et al.,
61 2014, 2016). Furthermore, cellular responses to *E. coli* O157 are associated with Th1
62 responses (Corbishley et al., 2014). The relationship between the shedding of *E. coli* and
63 immunity is not fully understood, but shedding has been associated with stressful events
64 that could affect the immune response (Ellis-Iversen et al., 2007; Munns et al., 2015).

65 Recent estimates using bulk milk antibody detection ELISAs based on fluke excretory-
66 secretory antigens show 50 - 80% of UK dairy herds have been exposed to fluke (McCann et
67 al., 2010; Howell et al., 2015). Although the current status for the beef sector is unknown,
68 figures released by the Food Standards Agency report that 16.5% of cattle livers were
69 condemned due to liver fluke during 2015 (Ford and Hadley, 2015). Since liver fluke

infection down-regulates Th1 responses, which are associated with clearance of the bacteria from the bovine gut (Corbishley et al., 2014), we hypothesized that fluke infection could affect the propensity of cattle to shed *E. coli* O157. If so, the presence of co-infected cattle could increase the risk of zoonotic *E. coli* infections.

2 Methods

This study was designed to utilise samples collected for an existing larger study on *E. coli* O157 in cattle intended for human consumption, funded by Food Standards Scotland (FSS) and the Food Standards Agency (FSA; Project FS101055); referred to below as the FSS/FSA study. For the FSS/FSA study, sample size calculations showed that a minimum of 110 Scottish farms and 160 farms from England and Wales were required to estimate a prevalence of *E. coli* O157 of 20% and 35% respectively within a tolerance of 0.168 with 95% confidence (Henry et al., 2017).

A sample size calculation to determine the number of cattle that were required to investigate the association between *F. hepatica* infection and *E. coli* O157 shedding was performed by Hickey et al. (2015) using simulated datasets. The estimated prevalence of *E. coli* O157 was set at 4% of cattle and 20% of farms (Pearce et al., 2009) whilst the estimated prevalence of *F. hepatica* was set at 20% of cattle and 80% of farms (Salimi-Bejestani et al., 2005b; McCann et al., 2010). 100% sensitivity and specificity of both tests were assumed. The result of using these parameters was that the inclusion of 1645 individual samples, from 50 randomly selected farms, would give the study a power of 87% to detect a two-fold increase in the odds that an animal would shed *E. coli* O157 if it was also infected with *F. hepatica*, compared to cattle not infected with fluke.

2.1 Sample and data collection

Two hundred and seventy farms were sampled in the FSS/FSA study (Henry et al., 2017). These included a variety of types of enterprise and breeds of cattle. Of these, 110 were

95 Scottish farms, randomly selected from all Scottish farms that had participated in both of
96 two earlier studies. The inclusion criterion was that there was at least one male aged one
97 year or over, or female over two without calves on the farm, as these farms were most
98 likely to contain animals that would end up in the food chain. In addition, 160 farms for
99 England and Wales were recruited from a randomly selected subset with either a male of
100 any breed aged over 1 year, or a female of a non-dairy breed aged over 1 year. Farmers
101 were initially notified by letter and given the choice to opt out, and were then contacted by
102 phone in a randomised order to enrol them in the study.

103 Farms were visited once between September 2014 and November 2015. Individual fresh
104 faecal pat samples were taken from the floor or ground, for the group of cattle from each
105 farm that contained the animals closest to going off the farm for slaughter. The number of
106 samples collected from each group was determined by a protocol assuming that if 8% of
107 animals were positive, there would be a 0.9 probability of identifying groups containing at
108 least one positive animal (Chase-Topping et al., 2007). It was assumed that a pat sample is
109 equivalent to an animal level unit for analysis. These samples were then sent to the
110 Epidemiology Research Unit (ERU) microbiological facilities at Scotland's Rural College
111 (SRUC), Inverness, within 48 hours of collection, and tested for *E. coli* O157. The
112 recruitment and visits were done by members of SRUC project team in Scotland, and the
113 ADAS project team in England and Wales.

114 Farms for which samples were submitted to SRUC's ERU laboratory on or after 5th January
115 2015, and which consented to further use of their samples and data for research purposes,
116 were included in the study. Delays due to funding and contractual issues meant that
117 samples received prior to this date were not retained.

118 Information on animal characteristics and farm management was collected from the
119 livestock keeper or farm manager on each farm, via a questionnaire administered by the

120 survey staff. The information was collected in an electronic format and was a shortened
121 version of a questionnaire used in a previous study (Chase-Topping et al., 2007). The
122 questionnaire was piloted with several farmers before use. The finalised questionnaire was
123 approved by the FSS Survey Control team (Henry et al., 2017). The questionnaire was
124 conducted in Welsh for Welsh-speaking respondents.

125 The information obtained was at the farm level, for example the age of animals was given
126 as a range for the group, and all animals in a group were treated as having been managed
127 the same in terms of housing, feeding and treatments given. The information relevant for
128 the current study was identified and extracted. As the aim was to develop a model to
129 determine the presence of an association between fluke and *E. coli* O157, rather than a
130 predictive model, only management information relevant to fluke was taken for use in the
131 model, to control for possible confounders which may be linked to both fluke and *E. coli*
132 O157. A summary of these is shown in Table 1.

133 **2.2 *E. coli* testing**

134 One gram of faeces was added to 20 ml of buffered peptone water (BPW, Thermo
135 Scientific, UK). The BPW was incubated for six hours at 37°C ($\pm 1^\circ$) then subjected to
136 immunomagnetic separation (IMS). Briefly, a 1 ml aliquot from each 20 ml BPW sample was
137 added to 20 μ l paramagnetic beads coated with polyclonal antibody for *E. coli* O157
138 lipopolysaccharide (Lab M Ltd., UK). The aliquots were mixed on a rotary mixer for 30
139 minutes before being washed three times in PBS with 0.05% Tween 20 (PBST, Sigma-Aldrich
140 Co. Ltd.). After the third wash, the beads were re-suspended in 100 μ l PBST and cultured
141 onto MacConkey agar containing sorbitol, cefixime (0.05 mg/l) and tellurite (2.5 mg/l) (CT-
142 SMac, Thermo Scientific, UK)(Jenkins et al., 2003).

143 Following overnight incubation at 37°C ($\pm 1^\circ$) plates were examined for non-sorbitol-
144 fermenting colonies and any suspect colonies were subcultured onto Chromocult coliform

145 agar (Merck KGaA., Germany). After a further overnight incubation at 37°C ($\pm 1^\circ$) any
146 resulting red colonies were tested with anti-*E. coli* O157 latex (Thermo Scientific, UK) for
147 agglutination. Colonies that agglutinated were identified as presumptively positive and
148 enumerated by limiting dilution.

149 Polymerase Chain Reaction (PCR) was used to confirm the serogroup of the isolates as
150 *E. coli* O157 (ISO/TS, 2012). For all positive samples, the number of *E. coli* O157 were
151 enumerated by culturing 10-fold dilutions of faeces in minimum recovery diluent, starting
152 from 1:10, on duplicate CT-SMac plates. Typical colonies were counted after overnight
153 incubation at 37°C ($\pm 1^\circ$) and counts expressed as colony forming units (cfu) per gram of
154 faeces.

155 IMS is considered to be a highly sensitive and specific method of identifying *E. coli* O157,
156 and has a lower limit of detection of 50 cfu/g (Aydin et al., 2014; Wright et al., 1994). Lower
157 cfu counts can be detected with decreased sensitivity. IMS has a specificity of 99% (Ekong
158 et al., 2017), and all positive isolates were confirmed as such by the Scottish *E. coli*
159 Reference Laboratory. For the positive/negative analysis, an *E. coli* O157 positive cow was
160 defined as one that tested positive by IMS. The limit of accurate enumeration was 100 cfu/g
161 of faeces (Pearce et al., 2004), and samples from which too few *E. coli* were cultured to be
162 enumerated were assigned a cfu/g of 10.

163 **2.3 *F. hepatica* testing**

164 Extraneous faecal material (2g), from each faecal sample was weighed into polypropylene
165 tubes and frozen (-20°C). When the *E. coli* O157 status of the farms was known (as defined
166 in Henry et al., 2017), all the samples from eligible *E. coli* O157 positive farms were
167 transported to Moredun Research Institute (MRI) in batches. Here they were tested using a
168 copro-antigen ELISA according to the manufacturer's instructions (Bio-X Diagnostics,
169 Jemelle, Belgium). MRI staff members were blinded to the *E. coli* O157 status of the

170 individual samples. Freezing the samples prior to performing the copro-antigen ELISA is
171 reported to make no difference to the sensitivity or specificity of the test (Brockwell et al.,
172 2013; Flanagan et al., 2011), and this was also confirmed before this study commenced
173 (Personal communication, Dr Philip Skuce).

174 The result was determined by calculating the percentage positivity (PP) of each sample
175 relative to the optical density (OD) of the positive control, after subtracting the OD of the
176 negative control (provided in the kit). The positive/negative cut off was determined by the
177 quality control insert supplied with the kit, and was either 7 or 8 for all the kits used for this
178 study. This test has a reported sensitivity of 0.77 and a specificity of 0.99 (Mazeri et al.,
179 2016).

180 **2.4 Statistical analysis**

181 The epidemiological unit of interest was the individual animal. For each animal for which a
182 sample was tested, the following results were obtained: *E. coli* O157 positive/negative, *E.*
183 *coli* O157 cfu/g, *F. hepatica* positive/negative derived by applying the cut off to the copro-
184 antigen ELISA results, and *F. hepatica* PP result (on a continuous scale). Farms without a
185 single fluke positive animal were excluded from further analysis, to ensure that cattle at
186 least had a possibility to be infected by fluke, which would not necessarily be the case if
187 there was no fluke on the farm. R (R Core Team, 2011) was used, with the lme4 (Bates et
188 al., 2015) and ggplot2 (Wickham, 2009) packages. Due to confidentiality agreements
189 relating to FSS/FSA project FS101055 which funded the faecal sample collection, figures or
190 data relating to groups of fewer than five farms cannot be shown.

191 **2.5 Multilevel model**

192 Correlations between the numerical explanatory variables were checked to ensure highly
193 correlated variables were not entered simultaneously into the model. All models were
194 fitted using maximum likelihood. Linear and logistic regression models were built with log₁₀
195 *E. coli* cfu/g and a positive *E. coli* result respectively as the outcome variable. Either log

196 fluke ELISA PP or a positive fluke result was used as the only level 1 explanatory variable,
197 and all other animal and farm management information were level 2 variables. One and 2
198 respectively were added to the *E. coli* O157 count and fluke ELISA PP results before logging
199 to deal with zero and negative values.

200 The starting point was a variable intercept model including a positive fluke result as a level
201 1 explanatory variable and farm as a level 2 random effect. Management variables which
202 met the inclusion criteria were then added one at a time. A seasonal pattern was expected
203 for *E. coli* (Ferens and Hovde, 2011), so day was modelled as a sinusoidal function to allow
204 for this. The same process was repeated with fluke PP as the explanatory variable. The
205 process was then repeated again with \log_{10} *E. coli* cfu/g as a continuous outcome variable.
206 Variable slopes were also tested. The Akaike information criterion (AIC) was used to
207 compare models, with a lower AIC considered better than a higher one.

208 **3 Results**

209 Between 13th January and 19th October 2015, of 39 farms sampled with one or more cattle
210 testing positive for *E. coli* O157, two declined to take part in further research and samples
211 from two farms were delayed in transit and were therefore not suitable for fluke testing.
212 There was insufficient sample for testing from a further five cows. Therefore, samples from
213 810 cattle from 35 herds were tested using the *F. hepatica* copro-antigen test. Of these, 14
214 farms had at least one cow testing positive for *F. hepatica*. Between 7 and 40 cattle were
215 sampled from each of these farms (median = 22, total = 334) and are included in the
216 following analysis.

217 **3.1 Descriptive statistics**

218 The characteristics of the farms are shown in Table 1. The data were examined to find out
219 whether groups of cattle were housed or grazing, how long they had been housed or
220 grazing for, and whether they had received a worming or flukicide treatment within the

221 past 3 months. However, even in groups for which flukicide use was recorded, fluke copro-
222 antigen ELISA positive cattle were still present, and similarly some groups of cattle which
223 had been housed for several months still had significant numbers of fluke positive animals.
224 Therefore treatment history was not used to exclude farms and all animals that came from
225 groups with at least one fluke case were included, on the basis that they would all have had
226 the chance to become fluke infected.

227 **3.1.1 Animal level**

228 Overall, 50.9% of cattle tested positive for *E. coli* O157 and 13.2% tested positive for *F.*
229 *hepatica*. The distributions are shown in figure 1.

230 **3.1.2 Farm level**

231 Within farms, between 4 and 100% of cattle tested positive for *E. coli* (mean = 43.5%,
232 median = 43.1%) whilst for *F. hepatica* the range was 2.1 to 100% (mean = 14.7%, median =
233 6.5%). The distribution of log *E. coli* O157 cfu/g varied between farms, but in general it was
234 right skewed in ten farms whilst four farms showed a more symmetrical platykurtic
235 distribution. For fluke PP, all except one farm had a right skewed distribution.

236 The farms were spread throughout Great Britain with six from Scotland, four from England
237 and four from Wales. North Wales, South Wales, the Welsh borders, Northern England and
238 a variety of Scottish locations were represented.

239 **3.2 Associations between fluke and *E. coli* O157**

240 Inspection of scatterplots revealed no visible association between the fluke PP and log *E.*
241 *coli* O157 cfu/g, either at individual or farm level (data not shown). More detailed
242 inspection of three farms with more than 10% of cattle testing positive for fluke revealed
243 no consistent pattern with regard to which individuals were positive for which pathogen. In
244 one farm all of the fluke positive animals were also *E. coli* positive, in a second farm all of
245 the fluke positives were *E. coli* negative, and in a third farm all animals had fluke and the PP
246 values were evenly spread between the *E. coli* positive and negative animals.

247 **3.3 Multi-level models**

248 The plotting of management variables against \log_{10} *E. coli* O157 cfu/g did not reveal any
249 non- linear relationships. No correlations of $r > 0.7$ were seen between any of the
250 explanatory variables, except between numbers of animals of different ages/types.

251 Four different combinations of output and input variables were tested, to include all
252 combinations of the *F. hepatica* and *E. coli* data. The inclusion of random effects improved
253 the model fit in every case, indicating that there were important differences between
254 farms. The explanatory management variables shown in Table 1 were each added to the
255 model as level 2 variables, but it was not possible to add more than two variables at once
256 because of the relatively small number of fluke cases, which led to non-convergence of the
257 model due to perfect partitioning.

258 The best models for each combination of *E. coli* O157 positive and \log_{10} *E. coli* O157 cfu/g,
259 and log fluke PP and fluke positive are shown (Table 2). The fluke result did not explain any
260 additional variation in three out of four models, however, log fluke PP was significant when
261 modelled against positive *E. coli* O157 result. Day of sampling and the age of the youngest
262 animal in the group were included in all of the models at level 2 and were highly significant
263 in all models ($p < 0.0001$). The higher the age of the youngest animal in the group, the
264 lower the odds of infection with *E. coli* O157. The model fitted better with day of sampling
265 as a linear variable, and the odds of *E. coli* O157 was found to decrease throughout the
266 year, from January until October. The introduction of random slopes worsened the model
267 fit in each case so this was not pursued.

268 **4 Discussion**

269 This study aimed to use samples available from the FSS/FSA study to investigate whether
270 shedding of *E. coli* O157 is associated with *F. hepatica* infection in cattle (Hickey et al.,
271 2015). *E. coli* O157 is the serogroup most commonly detected in humans in the UK, Europe

272 and the US, and is associated severe clinical outcomes in humans (Anon, 2015; Browning et
273 al., 2016). The advantage of using the samples from a pre-existing study was efficiency in
274 terms of reducing resources needed for planning, recruitment of farmers, visiting farms and
275 testing samples for *E. coli* O157. However, the biggest disadvantage of using the samples
276 gathered for the FSS/FSA study was that the sampling method was designed to treat the
277 group of cattle as the unit of interest; specifically, to identify groups where at least one
278 animal was shedding *E. coli* O157 (Gunn et al., 2007; Pearce et al., 2009). *E. coli* O157
279 shedding varies widely from day to day (Robinson et al., 2009), and the effective sensitivity
280 may be as low as 40% for a one-off faecal sample (Echeverry et al., 2005). Whilst this was
281 not a problem for the FSS/FSA study, where the group was treated as the epidemiological
282 unit, it may have affected the current study because individuals that were shedding *E. coli*
283 O157 may have been missed.

284 Reaching the required sample size depended on the initial assumptions about prevalence of
285 the two pathogens being reasonably accurate, particularly as the collection of additional
286 samples was not possible given the constraints of the study. However, the levels of fluke
287 infection seen in this study, both at the farm and the animal level, were much lower than
288 had been assumed for the sample size calculations (Hickey et al., 2015), and *F. hepatica*
289 only occurred in 43% of farms compared to the predicted 70-80% (Salimi-Bejestani et al.,
290 2005a; McCann et al., 2010; Hickey et al., 2015). This lower fluke prevalence may be partly
291 explained by differences between the cattle populations that were the subject of the
292 FSS/FSA study and previous studies. Data on herd level prevalence of infection is from
293 lactating dairy cows, whereas the FSS/FSA study sampled mostly beef breed or cross bred
294 store or finishing animals. Differences in management exist between these groups that are
295 likely to affect their risk of fluke infection. For example, treatment for fluke is more difficult
296 in dairy animals due to the long milk withhold times of flukicides. Also, of the 35 groups of
297 cattle tested for fluke, only nine were currently grazing, and of those, three had been

298 turned out onto pasture within the three weeks prior to sample collection. This could have
299 been due to the time of year when the samples were collected but also the nature of the
300 farming units tested. It is possible that some of the groups were permanently housed,
301 which would put them at low risk of fluke exposure, although this information was not
302 available from the questionnaire.

303 The lower than expected prevalence of fluke could also be due to the relatively low
304 sensitivity of the *F. hepatica* copro-antigen test, which in naturally infected cattle has been
305 estimated to be below 50%-60% (Duscher et al., 2011), whereas the bulk milk tank antibody
306 ELISA used to estimate prevalence in previous studies has a sensitivity of 96% (Salimi-
307 Bejestani et al., 2005a). The difference in sensitivity between the two types of test was not
308 taken into consideration in the feasibility study (Hickey et al., 2015). The relatively low
309 sensitivity of the diagnostic tests used for both fluke and *E. coli* could have led to non-
310 differential misclassification. This is expected to bias the observed effect size towards zero,
311 although sometimes, by chance, the effect size can be over-estimated (Jurek et al., 2005).

312 Delays in implementation of this study led to the loss of samples from 17 farms enrolled in
313 the FSS/FSA study that were sampled between September 2014 and January 2015 and
314 which had agreed to take part. This contributed to the failure to reach the required sample
315 size. The missed samples were taken during the season when the within-herd prevalence of
316 fluke might have been expected to be at its highest (Bloemhoff et al., 2015).

317 In spite of these challenges, one of our models showed a significant association between *F.*
318 *hepatica* and *E. coli* O157. This would be consistent with our initial hypothesis that *F.*
319 *hepatica* mediated down-regulation of Th-1 immunity may limit the ability of cattle to clear
320 *E. coli* O157 from the intestinal tract: indeed, this would be similar to the previous
321 observation that *F. hepatica* infections in cattle result in increased susceptibility to

322 *Salmonella dublin* which is associated with reduced cellular immune responses against the
323 bacteria (Aitken et al., 1979).

324 The cut-off of the fluke copro antigen ELISA has been the subject of debate, with some
325 studies setting their own cut-off to increase sensitivity (Brockwell et al., 2014). A
326 continuous measure of PP avoids this problem and PP is a biologically meaningful measure
327 as antigen level is correlated with fluke burden (Kamaludeen, 2016).

328 The addition of more than two additional explanatory variables was prevented by
329 insufficient variability within the data. This could partly explain the observed large random
330 effect of farm, which indicates that there were large differences between farms. Another
331 interesting question is whether the inter farm differences could be partially explained by
332 differences between strains of *E. coli* O157 at the molecular level that might be related to
333 shedding events and immune status. Indeed it is known that different strains of *E. coli* O157
334 induce different types of immune response (Corbishley et al., 2014) and different genetic
335 traits of *E. coli* O157, such as phage type (Chase-Topping et al., 2007), presence of *stx2a*
336 and *stx2c* genes and polymorphisms in the *tir* gene (Arthur et al., 2013), are associated with
337 either high or low shedding from infected individuals. Although it was important to control
338 for day of sampling, as season is associated with observed prevalence of both fluke and *E.*
339 *coli* O157 (Bloemhoff et al., 2015; Ferens and Hovde, 2011; Smith et al., 2016; Synge et al.,
340 2003), the strong effect seen here is more likely to be due to all animals from a single farm
341 being sampled on the same day than a genuine seasonal effect. Therefore caution should
342 be used when interpreting the direction and size of the seasonal effect. There may also be
343 other explanatory or confounding variables that are not included in these models. The
344 result should therefore be interpreted with caution, particularly as the effect size is small
345 and it is only seen in one of the model combinations.

346 The results of our study hint at an association between *E. coli* O157 shedding and *F.*
347 *hepatica* infection that merits further investigation. Based on our experiences, use of pre-
348 collected samples represents a cost-effective way of obtaining data, however, care needs
349 to be taken to avoid certain pit falls. In the planning stage of future studies, worst case
350 scenarios for prevalence should be considered, taking into account diagnostic test accuracy
351 and differences between populations which may affect apparent prevalence. Even more
352 importantly, efforts should be concentrated on ensuring that the true infection or shedding
353 status of each individual can be ascertained, and that the type, number and size of sample
354 are suitable for this. For *E. coli* shedding this is likely to include longitudinal sampling to
355 address the issues of intermittent shedding and uneven distribution of bacteria within the
356 faeces.

357 **Acknowledgements**

358 The funding source for the collection of the original samples was the FSS and FSA, project
359 number FS101055: *E. coli* O157 super-shedding in cattle and the mitigation of human risk.
360 Copro-antigen testing was funded by the Moredun Foundation. AH was funded by a BBSRC
361 doctoral training partnership (DTP). Conflict of interest: None.

362 We are grateful for the contribution of the SRUC ERU and ADAS project team members and
363 thank those livestock keepers who gave permission for the additional use of their data and
364 the faecal samples from their holdings for further research.

365 **References**

366 Aitken, M.M., Hughes, D.L., Jones, P.W., Hall, G.A., Smith, G.S., 1979. Immunological
367 responses of fluke-infected and fluke-free cattle to *Salmonella dublin* and other
368 antigens. Res. Vet. Sci. 27, 306–12.

369 Aitken, M.M., Jones, P.W., Hall, G.A., Hughes, D.L., Brown, G.T., 1981. Responses of fluke-

370 infected and fluke-free cattle to experimental reinfection with *Salmonella dublin*. Res.
371 Vet. Sci. 31, 120–6.

372 Anon, 2015. Annual epidemiological report on Shigatoxin/verocytotoxin-producing
373 *Escherichia coli* (STEC/VTEC) infection. Stockholm.

374 Aydin, M., Herzig, G.P.D., Jeong, K.C., Dunigan, S., Shah, P., Ahn, S., 2014. Rapid and
375 sensitive detection of *Escherichia coli* O157:H7 in milk and ground beef using
376 magnetic bead-based immunoassay coupled with tyramide signal amplification. J.
377 Food Prot. 77, 100–5. doi:10.4315/0362-028X.JFP-13-274

378 Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models
379 Using lme4. J. Stat. Softw. 67. doi:10.18637/jss.v067.i01

380 Bloemhoff, Y., Forbes, A., Danaher, M., Good, B., Morgan, E., Mulcahy, G., Sekiya, M.,
381 Sayers, R., 2015. Determining the Prevalence and Seasonality of *Fasciola hepatica* in
382 Pasture-based Dairy herds in Ireland using a Bulk Tank Milk ELISA. Ir. Vet. J. 68, 16.
383 doi:10.1186/s13620-015-0042-5

384 Brady, M.T., O'Neill, S.M., Dalton, J.P., Mills, K.H.G., 1999. *Fasciola hepatica* suppresses a
385 protective Th1 response against *Bordetella pertussis*. Infect. Immun. 67, 5372–5378.

386 Brockwell, Y.M., Elliott, T.P., Anderson, G.R., Stanton, R., Spithill, T.W., Sangster, N.C., 2014.
387 Confirmation of *Fasciola hepatica* resistant to triclabendazole in naturally infected
388 Australian beef and dairy cattle. Int. J. Parasitol. Drugs Drug Resist. 4, 48–54.
389 doi:10.1016/j.ijpddr.2013.11.005

390 Brockwell, Y.M., Spithill, T.W., Anderson, G.R., Grillo, V., Sangster, N.C., 2013. Comparative
391 kinetics of serological and coproantigen ELISA and faecal egg count in cattle
392 experimentally infected with *Fasciola hepatica* and following treatment with

393 triclabendazole. *Vet. Parasitol.* 196, 417–426. doi:10.1016/j.vetpar.2013.04.012

394 Browning, L., Allison, L., Couper, S., Hanson, M., Hawkins, G., Smith-Palmer, A., 2016. STEC
395 in Scotland 2016: enhanced surveillance and reference laboratory data. *Heal. Prot.*
396 *Scotl. Wkly. Rep.* 51.

397 Chase-Topping, M.E., McKendrick, I.J., Pearce, M.C., MacDonald, P., Matthews, L., Halliday,
398 J., Allison, L., Fenlon, D., Low, J.C., Gunn, G., Woolhouse, M.E.J., 2007. Risk factors for
399 the presence of high-level shedders of *Escherichia coli* O157 on Scottish farms. *J. Clin.*
400 *Microbiol.* 45, 1594–1603. doi:10.1128/JCM.01690-06

401 Corbishley, A., Ahmad, N.I., Hughes, K., Hutchings, M.R., McAteer, S.P., Connelley, T.K.,
402 Brown, H., Gally, D.L., McNeilly, T.N., 2014. Strain-dependent cellular immune
403 responses in cattle following *Escherichia coli* O157:H7 colonization. *Infect. Immun.* 82,
404 5117–31. doi:10.1128/IAI.02462-14

405 Corbishley, A., Connelley, T., Wolfson, E., Ballingall, K., Beckett, A., Gally, D., McNeilly, T.,
406 2016. Identification of epitopes recognised by mucosal CD4+ T-cell populations
407 from cattle experimentally colonised with *Escherichia coli* O157:H7. *Vet. Res.* 47.

408 DEFRA, 2005. Report on TB SE3013.

409 Duscher, R., Duscher, G., Hofer, J., Tichy, A., Prosl, H., Joachim, A., 2011. *Fasciola hepatica* -
410 Monitoring the milky way? The use of tank milk for liver fluke monitoring in dairy
411 herds as base for treatment strategies. *Vet. Parasitol.* 178, 273–278.
412 doi:10.1016/j.vetpar.2011.01.040

413 Echeverry, A., Loneragan, G.H., Wagner, B.A., Brashears, M.M., 2005. Effect of intensity of
414 fecal pat sampling on estimates of *Escherichia coli* O157 prevalence. *Am. J. Vet. Res.*
415 66, 2023–7. doi:10.2460/ajvr.2005.66.2023

416 Ekong, P.S., Sanderson, M.W., Bello, N.M., Noll, L.W., Cernicchiaro, N., Renter, D.G., Bai, J.,
 417 Nagaraja, T.G., 2017. Bayesian estimation of true prevalence, sensitivity and specificity
 418 of three diagnostic tests for detection of *Escherichia coli* O157 in cattle feces. *Prev.*
 419 *Vet. Med.* 148, 21–27. doi:10.1016/J.PREVETMED.2017.10.002

420 Ellis-Iversen, J., Smith, R.P., Snow, L.C., Watson, E., Millar, M.F., Pritchard, G.C., Sayers, A.R.,
 421 Cook, A.J.C., Evans, S.J., Paiba, G.A., 2007. Identification of management risk factors
 422 for VTEC O157 in young-stock in England and Wales. *Prev. Vet. Med.* 82, 29–41.
 423 doi:10.1016/j.prevetmed.2007.05.004

424 Ferens, W.A., Hovde, C.J., 2011. *Escherichia coli* O157:H7: Animal Reservoir and Sources of
 425 Human Infection. *Foodborne Pathog. Dis.* 8, 465–487.

426 Flanagan, A., Edgar, H., Forster, F., Gordon, A., Hanna, R., McCoy, M., Brennan, G.,
 427 Fairweather, I., 2011. Standardisation of a coproantigen reduction test (CRT) protocol
 428 for the diagnosis of resistance to triclabendazole in *Fasciola hepatica*. *Vet. Parasitol.*
 429 176, 34–42. doi:10.1016/J.VETPAR.2010.10.037

430 Flynn, R.J., Mannion, C., Golden, O., Hacariz, O., Mulcahy, G., 2007. Experimental *Fasciola*
 431 *hepatica* Infection Alters Responses to Tests Used for Diagnosis of Bovine
 432 Tuberculosis. *Infect. Immun.* 75, 1373–1381. doi:10.1128/IAI.01445-06

433 Flynn, R.J., Mulcahy, G., Elsheikha, H.M., 2010. Coordinating innate and adaptive immunity
 434 in *Fasciola hepatica* infection: Implications for control. *Vet. Parasitol.* 169, 235–240.
 435 doi:10.1016/j.vetpar.2010.02.015

436 Ford, F., Hadley, P., 2015. Minimising carcase losses for Better Returns BEEF AND SHEEP
 437 BRP MANuAl 9.

438 Garza-Cuartero, L., Blanco, A., McNair, J., Flynn, R.J., Williams, D., Diggle, P., Cassidy, J.,

439 Mulcahy, G., 2016. *Fasciola hepatica* Infection Reduces *Mycobacterium* and
 440 *Mycobacterial* Uptake Suppresses the Pro-inflammatory Response. *Parasite Immunol.*
 441 38, 387–402. doi:10.1111/pim.12326

442 Graham-Brown, J., 2016. Immune responses in dairy cattle naturally exposed to *fasciola*
 443 *hepatica*. University of Liverpool.

444 Gunn, G.J., McKendrick, I.J., Ternent, H.E., Thomson-Carter, F., Foster, G., Synge, B.A., 2007.
 445 An investigation of factors associated with the prevalence of verocytotoxin producing
 446 *Escherichia coli* O157 shedding in Scottish beef cattle. *Vet. J.* 174, 554–564.
 447 doi:10.1016/j.tvjl.2007.08.024

448 Gyles, C.L., 2007. Shiga toxin-producing *Escherichia coli*: an overview. *J. Anim. Sci.* 85.
 449 doi:10.2527/jas.2006-508

450 Health Protection Network, 2013. Guidance for the Public Health Management of Infection
 451 with Verotoxigenic *Escherichia coli* (VTEC).

452 Henry, M.K., Tongue, S.C., Evans, J., Webster, C., McKendrick, I.J.I., Morgan, M., Willet, A.,
 453 Reeves, A., Humphry, R.W.R., Gally, D.L.D., Gunn, G.J.G., Chase-Topping, M.E.M.,
 454 Willett, A., Reeves, A., Humphry, R.W.R., Gally, D.L.D., Gunn, G.J.G., Chase-Topping,
 455 M.E.M., 2017. British *E. coli* O157 in Cattle Study (BECS): to determine the prevalence
 456 of *Escherichia coli* O157 in herds with cattle destined for the food chain. *Epidemiol.*
 457 *Infect.* 1–12. doi:10.1017/S0950268817002151

458 Hickey, G.L., Diggle, P.J., McNeilly, T.N., Tongue, S.C., Chase-Topping, M.E., Williams, D.J.L.,
 459 2015. The feasibility of testing whether *Fasciola hepatica* is associated with increased
 460 risk of verocytotoxin producing *Escherichia coli* O157 from an existing study protocol.
 461 *Prev. Vet. Med.* doi:10.1016/j.prevetmed.2015.02.022

462 Howell, A., Baylis, M., Smith, R., Pinchbeck, G., Williams, D., 2015. Epidemiology and impact
 463 of *Fasciola hepatica* exposure in high-yielding dairy herds. *Prev. Vet. Med.* 121, 41–48.
 464 doi:10.1016/j.prevetmed.2015.05.013

465 Jenkins, C., Pearce, M.C., Smith, A.W., Knight, H.I., Shaw, D.J., Cheasty, T., Foster, G., Gunn,
 466 G.J., Dougan, G., Smith, H.R., Frankel, G., 2003. Detection of *Escherichia coli*
 467 serogroups O26, O103, O111 and O145 from bovine faeces using immunomagnetic
 468 separation and PCR/DNA probe techniques. *Lett. Appl. Microbiol.* 37, 207–12.

469 Jurek, A.M., Greenland, S., Maldonado, G., Church, T.R., 2005. Proper interpretation of non-
 470 differential misclassification effects: expectations vs observations. *Int. J. Epidemiol.*
 471 34, 680–687. doi:10.1093/ije/dyi060

472 Kamaludeen, J., 2016. The Detection and Occurrence of Anthelmintic Resistance in *Fasciola*
 473 *hepatica* in Sheep in the United Kingdom. University of Liverpool.

474 Locking, M.E., O'Brien, S.J., Reilly, W.J., Wright, E.M., Campbell, D.M., Coia, J.E., Browning,
 475 L.M., Ramsay, C.N., 2001. Risk factors for sporadic cases of *Escherichia coli* O157
 476 infection: the importance of contact with animal excreta. *Epidemiol. Infect.* 127, 215–
 477 20.

478 Maizels, R.M., Balic, A., Gomez-Escobar, N., Nair, M., Taylor, M.D., Allen, J.E., 2004.
 479 Helminth parasites - Masters of regulation. *Immunol. Rev.* 201, 89–116.
 480 doi:10.1111/j.0105-2896.2004.00191.x

481 Mazeri, S., Sargison, N., Kelly, R.F., Bronsvoort, B.M. deC., Handel, I., Brennan, G., 2016.
 482 Evaluation of the performance of five diagnostic tests for *Fasciola hepatica* infection in
 483 naturally infected cattle using a Bayesian no gold standard approach. *PLoS One* 11,
 484 e0161621. doi:10.1371/journal.pone.0161621

485 McCann, C.M., Baylis, M., Williams, D.J.L., 2010. Seroprevalence and spatial distribution of
 486 *Fasciola hepatica*-infected dairy herds in England and Wales. *Vet. Rec.* 166, 612–617.
 487 doi:10.1136/vr.b4836

488 Moreau, E., Chauvin, A., 2010. Immunity against Helminths: Interactions with the Host and
 489 the Intercurrent Infections. *J. Biomed. Biotechnol.* doi:10.1155/2010/428593

490 Munns, K.D., Selinger, L.B., Stanford, K., Guan, L., Callaway, T.R., McAllister, T.A., 2015.
 491 Perspectives on Super-Shedding of *Escherichia coli* O157:H7 by Cattle. *Foodborne*
 492 *Pathog. Dis.* 12, 89–103. doi:10.1089/fpd.2014.1829

493 Paiba, G.A., Wilesmith, J.W., Evans, S.J., Pascoe, S.J.S., Smith, R.P., Kidd, S.A., Ryan, J.B.M.,
 494 McLaren, I.M., Chappell, S.A., Willshaw, G.A., Cheasty, T., French, N.P., Jones, T.W.H.,
 495 Buchanan, H.F., Challoner, D.J., Colloff, A.D., Cranwell, M.P., Daniel, R.G., Davies, I.H.,
 496 Duff, J.P., Hogg, R.A.T., Kirby, F.D., Millar, M.F., Monies, R.J., Nicholls, M.J., Payne, J.H.,
 497 2003. Prevalence of faecal excretion of verocytotoxigenic *Escherichia coli* O157 in
 498 cattle in England and Wales. *Vet. Rec.* 153, 347–353. doi:10.1136/vr.153.12.347

499 Pearce, M.C., Chase-Topping, M.E., McKendrick, I.J., Mellor, D.J., Locking, M.E., Allison, L.,
 500 Ternent, H.E., Matthews, L., Knight, H.I., Smith, A.W., Synge, B. a, Reilly, W., Low, J.C.,
 501 Reid, S.W.J., Gunn, G.J., Woolhouse, M.E.J., 2009. Temporal and spatial patterns of
 502 bovine *Escherichia coli* O157 prevalence and comparison of temporal changes in the
 503 patterns of phage types associated with bovine shedding and human *E. coli* O157
 504 cases in Scotland between 1998-2000 and 2002-2004. *BMC Microbiol.* 9, 276.
 505 doi:10.1186/1471-2180-9-276

506 Pearce, M.C., Fenlon, D., Low, J.C., Smith, A.W., Knight, H.I., Evans, J., Foster, G., Synge,
 507 B.A., Gunn, G.J., 2004. Distribution of *Escherichia coli* O157 in bovine fecal pats and its
 508 impact on estimates of the prevalence of fecal shedding. *Appl. Environ. Microbiol.* 70,

509 5737–5743. doi:10.1128/AEM.70.10.5737-5743.2004

510 Pritchard, G.C., Forbes, A.B., Williams, D.J.L., Salimi-Bejestani, M.R., Daniel, R.G., 2005.

511 Emergence of fasciolosis in cattle in East Anglia. *Vet. Rec.* 157, 578–582.

512 Pruimboom-Brees, I.M., Morgan, T.W., Ackermann, M.R., Nystrom, E.D., Samuel, J.E.,

513 Cornick, N. a, Moon, H.W., 2000. Cattle lack vascular receptors for *Escherichia coli*

514 O157:H7 Shiga toxins. *Proc. Natl. Acad. Sci. U. S. A.* 97, 10325–10329.

515 doi:10.1073/pnas.190329997

516 Public Health England, 2013. *Escherichia coli* (*E. coli*) O157 annual totals in England and

517 Wales [WWW Document]. URL

518 [https://www.gov.uk/government/publications/escherichia-coli-e-coli-o157-annual-](https://www.gov.uk/government/publications/escherichia-coli-e-coli-o157-annual-totals/escherichia-coli-e-coli-o157-annual-totals-in-england-and-wales)

519 [totals/escherichia-coli-e-coli-o157-annual-totals-in-england-and-wales](https://www.gov.uk/government/publications/escherichia-coli-e-coli-o157-annual-totals/escherichia-coli-e-coli-o157-annual-totals-in-england-and-wales) (accessed

520 11.8.16).

521 R Core Team, 2011. R: A language and environment for statistical computing. R Foundation

522 for Statistical Computing.

523 Robinson, S.E., Brown, P.E., Wright, E.J., Hart, C.A., French, N.P., 2009. Quantifying within-

524 and between-animal variation and uncertainty associated with counts of *Escherichia*

525 *coli* O157 occurring in naturally infected cattle faeces. *J. R. Soc. Interface* 6, 169–77.

526 doi:10.1098/rsif.2008.0183

527 Salimi-Bejestani, M.R., Daniel, R.G., Felstead, S.M., Cripps, P.J., Mahmood, H., Williams,

528 D.J.L., 2005a. Prevalence of *Fasciola hepatica* in dairy herds in England and Wales

529 measured with an ELISA applied to bulk-tank milk. *Vet. Rec.* 156, 729–731.

530 Salimi-Bejestani, M.R., McGarry, J.W., Felstead, S., Ortiz, P., Akca, A., Williams, D.J.L., 2005b.

531 Development of an antibody-detection ELISA for *Fasciola hepatica* and its evaluation

532 against a commercially available test. *Res. Vet. Sci.* 78, 177–181.
 533 doi:10.1016/j.rvsc.2004.08.005

534 Smith, R.P., Pollitt, W.J., Paiba, G.A., 2016. A longitudinal study of risk factors for shedding
 535 of VTEC O157 by young cattle in herds with known *E. coli* O157 carriage. *Epidemiol.*
 536 *Infect.* 144, 1818–1829. doi:10.1017/S095026881600008X

537 Strachan, N.J.C., Dunn, G.M., Locking, M.E., Reid, T.M.S., Ogden, I.D., 2006. *Escherichia coli*
 538 O157: Burger bug or environmental pathogen? *Int. J. Food Microbiol.* 112, 129–137.
 539 doi:10.1016/j.ijfoodmicro.2006.06.021

540 Synge, B.A., Chase-Topping, M.E., Hopkins, G.F., McKendrick, I.J., Thomson-Carter, F., Gray,
 541 D., Rusbridge, S.M., Munro, F.I., Foster, G., Gunn, G.J., 2003. Factors influencing the
 542 shedding of verocytotoxin-producing *Escherichia coli* O157 by beef suckler cows.
 543 *Epidemiol. Infect.* 130, 301–12.

544 Wickham, H., 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

545 Wright, D.J., Chapman, P.A., Siddons, C.A., 1994. Immunomagnetic separation as a sensitive
 546 method for isolating *Escherichia coli* O157 from food samples. *Epidemiol. Infect.* 113,
 547 31–39. doi:10.1017/S0950268800051438

548

549

550 **Table 1. Characteristics of the animals and farms in the fluke and *E. coli* O157 study**

| | Farm-level (n=14) | Individual animals (n=334) |
|---|--------------------------------|----------------------------|
| Day of sample collection (Day 1=1st Jan 2014) | Range = 20-293 Median = 126 | |
| Grazing | 3 (21.43 %) | |
| Housed | 11 (78.57 %) | |
| Dairy | 2 (14.29 %) | |
| Suckler beef | 8 (57.14 %) | |
| Finisher | 2 (14.29 %) | |
| Other | 2 (14.29 %) | |
| Youngest in group (months) | Range = 6-26 Median = 14.5 | |
| Oldest in group | Range = 11-48 Median = 20 | |
| Total number of cattle on farm | Range = 41-516 Median = 117 | |
| Total number of cows on farm (Females that have had a calf) | Range = 0-208 Median = 33 | |
| Total number of heifers on farm | Range = 0-65 Median = 6 | |
| Total number of cattle under 1 year on farm | Range = 0-215 Median = 30 | |
| Total number of ewes on farm | Range = 0-700 Median = 0 | |
| Total number of sheep overwintering on farm | Range = 0-433 Median = 0 | |
| Water supply from mains | 10 (71.43 %) | |
| Water supply from spring or well | 6 (42.86 %) | |
| Water supply from natural source | 11 (78.57 %) | |
| Median percentage of fluke positive cows ¹ | 6.55% 2.13-100% | |
| Range of positive cows | | |
| Fluke positive | | 44 (13.17%) |
| Median PP | | 0.82 |
| Range PP | | -1.07-73.74 |
| Median percentage of <i>E. coli</i> O157 positive cows ² | 43.10% | |
| Range of positive cows | 4.00-100% | |
| <i>E. coli</i> positive | | 170 (50.9%) |
| Median cfu/g ³ | | 10 |
| Range cfu/g ³ | | 0-1.45 x 10 ⁵ |

551 ¹'Fluke positive' refers to an animal which tested positive on the copro-antigen ELISA result

552 ²'*E. coli* O157 positive' refers to an animal with a positive IMS *E. coli* test

553 ³Samples from which *E. coli* numbers fell below the limit of enumeration were assigned cfu/g = 10




Figure 1. Distribution of *F. hepatica* coproantigen PP values and *E. coli* O157 cfu/g for animals across all farms.

Table 2. Summary of the multi-level models. Farm was included as a random intercept. Day of sampling and age of the youngest animal in the group were controlled for in all models.

| Outcome variable | Input variable | Co-efficient | <i>p</i> value |
|---------------------------------------|---------------------------------|--------------|----------------|
| <i>E. coli</i> O157 positive | <i>F. hepatica</i> positive | 0.50 | 0.34 |
| <i>E. coli</i> O157 positive | Log <i>F. hepatica</i> ELISA PP | 0.48 | 0.010 |
| log ₁₀ <i>E. coli</i> O157 | <i>F. hepatica</i> positive | -0.02 | 0.90 |
| log ₁₀ <i>E. coli</i> O157 | Log <i>F. hepatica</i> ELISA PP | 0.09 | 0.26 |

E. coli O157 positivity was determined using immune-magnetic separation. log₁₀ *E. coli* O157 refers log₁₀ of the *E. coli* O157 count (cfu/g). *F. hepatica* positivity was determined using a copro-antigen ELISA. ELISA PP is the percentage positivity compared to a known positive sample.